

Non-Intravenous Conscious Sedation Consent Form

Oral Sedation is provided to minimize anxiety and reduce restlessness during extensive dental procedures. It is safe, effective and its effect diminishes soon after the procedure has been completed. It is imperative however that the protocol described in this document be closely followed.

1. Please read and ask any questions, understand and sign this form and your surgical consent form prior to taking your pre-operative medication. You are unable to give informed consent while medicated and your procedure will not be performed without your valid, informed consent.
2. The medication you have been prescribed is a short acting drug in the benzodiazepine class. This medication has mild sedative and muscle relaxing properties. Dosage instructions will be given by your doctor, written on the bottle and should be followed exactly.
3. Because this medication is mildly sedating, it effects judgment and response time, and will make you sleepy. Therefore use of this medication in conjunction with dental treatment requires you to be escorted to and from your appointment. You should arrange a responsible escort and confirm with them well in advance of your appointment. Your escort must remain in the vicinity of the dental clinic for the duration of the appointment.
4. Because the effect of this medication continues for an indeterminate period after treatment, driving or operating of any heavy equipment is prohibited for a period of 12 hours after administration of this medication.
5. Oral Sedatives should not be ingested if you are pregnant, or breast-feeding. They should not be taken if you have kidney or liver disease, if you are hypersensitive to benzodiazepines (Valium, Ativan, Versed, Halcion, Xanax etc) or are taking any medication which adversely interact or amplify the effects of these medications (see medication reconciliation).
6. It is not necessary to fast prior to dental treatment with Oral Conscious Sedation; in fact a light meal or nutritious shake is encouraged. After your procedure you will remain mildly sedated and should rest, preferably with a companion present, as you may be unsteady and could become disoriented.

By checking the box that follows, you acknowledge that you read, speak and communicate English and that you have received adequate information about the treatment, you understand this information, you're willing to abide by the conditions described in this document, and that all

of your questions have been answered fully.